Written on OCTOBER 1, 2014 AT 4:13 PM by SVANDERWERFF

Controlling Your Blood Sugar During Pregnancy

Filed under FORCE HEALTH AND SAFETY, HEATH, PUBLIC HEALTH

(ONE COMMENT)

By Kelly O. Elmore, MD, MBA, Board Certified OBGYN

Gestational Diabetes

Navy Medicine is a global healthcare network of 63,000 Navy medical personnel around the world who provide high quality health care to more than one million eligible beneficiaries. Navy Medicine personnel deploy with Sailors and Marines worldwide, providing critical mission support aboard ship, in the air, under the sea and on the battlefield.

If you have been diagnosed with diabetes prior to pregnancy this is called Pre-gestat

I got a text from a friend today, "My 1 hour sugar test is elevated. What does that mean?" After we discussed expectations, the next week she texts again, "my 3 hour test is normal, do I need to do anything different for the rest of my pregnancy?"

Although there are many great books and websites out there to tell you what to expect during pregnancy, until it happens to you- sometimes you do not get to that chapter. Then once you get to your appointment, it may be all about logistics and you miss 85% of the conversation your provider had with you.

Here's what you need to know- short and sweet so you can remember. Don't worry I'll give you references at the end too.

Diabetes occurs when your blood sugar level in the blood is not Navy Medicine Social Media

Navy Medicine Video

Follow us on Twitter

facebook Join us on Facebook

issuv Read our publications

flickr View our photo stream

You Tube W

Watch our videos

Navy Medicine Live Archives

February 2015 (2)

January 2015 (12)

December 2014 (17)

November 2014 (11)

regulated. If you have been diagnosed with diabetes prior to pregnancy this is called **Pregestational**

diabetes. It is crucial to have this controlled prior to pregnancy to reduce major defects and risk of death in the early



Diabetes occurs when your blood sugar level in the blood is not regulated.

growth phases of your baby. If you are diagnosed with diabetes during your pregnancy then you have gestational (during pregnancy) diabetes. If it is very early in your pregnancy then you probably had it prior to pregnancy too.

Why did this happen to me? You naturally make a hormone called insulin that regulates your blood sugar levels. It is thought that the placenta carries hormones that alter your body's ability to stay balanced in order to provide nutrients to your baby.1 Some people's bodies respond well to this and others don't, especially if you are at higher risk prior to pregnancy.

When and how do I get diagnosed? Typically, between 24-28 weeks of pregnancy, your provider will ask you to drink a sugary drink (you don't have to fast) and wait one hour for a blood test. If you have risk factors: over the age of 30, African American, Hispanic, Native American heritage, elevated BMI (body mass index based on your height and weight), prior history of abnormal glucose testing, prior history of a large baby at delivery, you may have to do this test twice- once in the early 2nd trimester and again in the late 2nd trimester. Depending on the scale that is used, if your 1hr test if >140 (some clinics use 135), then you are at risk for GDM and a 2nd confirmation test will need to be done. For this test, you should fast for at least 8 hours. You will get blood drawn 4 times, once fasting (prior to eating), then you will drink more of the sugary drink, then drawn again at 1, 2, and 3hrs after drinking. Some people do vomit and have to try again another day or do an alternative method of testing, but most women tolerate the test very well. Once again depending on the scale used if two or more of the blood tests are abnormal or one blood level is significantly high, you are diagnosed with Gestational Diabetes.



If my 1 hour is abnormal and my 3 hour is normal am I free and clear? Absolutely not. Although you do not have to go through the steps explained below, research shows that you could potentially have insulin resistance and like

	ober 2014 (15) tember 2014 (20)
	Fust 2014 (14)
	7 2014 (13)
	e 2014 (8)
	y 2014 (11)
	il 2014 (9)
	rch 2014 (14)
	ruary 2014 (7)
	uary 2014 (7)
	ember 2013 (7)
	vember 2013 (12)
	ober 2013 (7)
Sep	tember 2013 (14)
Aug	tust 2013 (13)
Jul	⁷ 2013 (11)
Jun	e 2013 (22)
Ma	y 2013 (15)
Apı	il 2013 (14)
Ma	rch 2013 (14)
Feb	ruary 2013 (14)
Jan	uary 2013 (12)
Dec	ember 2012 (11)
Nov	vember 2012 (11)
Oct	ober 2012 (7)
Sep	tember 2012 (9)
Aug	just 2012 (12)
Jul	2012 (13)
Jun	e 2012 (17)
Ma	y 2012 (22)
Apı	il 2012 (14)
Ma	rch 2012 (13)
Feb	ruary 2012 (14)
	uary 2012 (13)
	ember 2011 (13)

Research shows that you could potentially have insulin resistance and like everyone during their pregnancy (barring other medical concerns) daily exercise of at least 30 minutes and monitoring your eating choices is very important.

everyone during their
pregnancy (barring
other medical
concerns) daily
exercise of at least 30
minutes and
monitoring your

October 2011 (22)	
September 2011 (12)	
August 2011 (16)	
July 2011 (10)	

eating choices is very important.

If my 3 hour is abnormal then what's next? Your provider should order you a set of supplies to check your blood sugar 4-6 times a day. Your provider will review goals of glucose level management, I recommend fasting glucose <95 and 2 hours after meals < 120 or <140 if 1 hour after a meal. If levels are not controlled within 2-3 weeks with diet and exercise then medication therapy is recommended. Medications may either be oral or through an injection.

Will I get delivered early? Do I need additional testing?

This is all dependent on how well our body responds to your change in exercise and eating habits. You may undergo fetal testing (monitoring the fetal heart rate and measuring the fluid levels around your baby) during the latter part of your 3rd trimester, you may need a growth ultrasound to help calculate the baby's weight, and/or delivery prior to your due date. But remember, the goal is to have a healthy baby so do your best and your provider will do the same to help you reach your goal.

I'm delivered, what next? If you are well controlled most providers stop checking your blood sugars after delivery but this does not mean you should change your healthy eating habits. Also, as soon are you are able to return to working out you should do so. By the way, breastfeeding is a great way to reduce you and your newborns risk for developing diabetes in the future. Around your 5-6th week postpartum, you will need to do another fasting sugar test but this one takes 2 hours. This is to make sure your



Always continue to eat healthy and exercise.

diabetes has resolved. If it has not, you will need to see your provider and continue with the nutritionist to get your levels controlled and help prevent major complications from diabetes in the future. You should also continue annual visits with your primary care provider and check at least every 2-3 years for the development of Type II Diabetes. You should also inform your baby's physician that you had diabetes during pregnancy because your child is at risk for childhood diabetes as well.

Always continue to eat healthy and exercise.


The above information is for educational purposes only. Please discuss in depth with your OB provider.

References:

ACOG bulletin FAQ 177 obtained September 2014

http://www.acog.org/Patients/FAQs/Gestational-Diabetes

American College of Obstetricians and Gynecologists Committee on Practice Bulletins—Obstetrics. ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists. Number 30, September 2001 (replaces Technical Bulletin Number 200, December 1994). Gestational diabetes. *Obstet Gynecol.* 2001;98(3):525-538.



http://www.labradordellaspippola.it/ Martino Porta interesting!

lab st